



Magneto-optics of massless electrons in Cd_3As_2

Milan Orlita

Laboratoire National des Champs Magnétiques Intenses
CNRS, Grenoble, France



Collaborators:

M. Hakl, C. Faugeras, B. A. Piot,
G. Martinez, M. Potemski
A. Akrap, I. Crassee, D. van der Marel
F. Teppe, W. Desrat
S. Tchoumakov, M. O. Goerbig
C. C. Homes
A. Arushanov, A. Nateprov
Q. D. Gibson, R. J. Cava
J. Kuba, O. Caha, J. Novák
S. Koohpayeh, L. Wu, N. P. Armitage

LNCMI, CNRS, Grenoble, France
Université de Genève, Switzerland
LCC-CNRS & Université Montpellier, France
LPS-CNRS, Paris Orsay, France
Brookhaven national Laboratory, USA
Institute of Applied Physics, ASM, Moldova
Princeton University, USA
CEITEC & Masaryk University, Brno, Czech Republic
Johns Hopkins University, Baltimore, USA




Outline:

- Massless electrons in solid-state physics (reminder)
- 3D massless electrons in Cd_3As_2
- Conclusions



Graphene

2D



2D crystal made of carbon atoms organized in hexagonal lattice

Theoretically known over
sixty years...

P. R. Wallace, Phys. Rev. 71, 622 (1947)

Isolated/fabricated in 2004/2005

K. S. Novoselov et al., Science 306, 666 (2004)

K. S. Novoselov et al., Nature 438, 197 (2005)

3D

Diamond




3D

Graphite




1D

Carbon
nanotube



Fullerenes




0D




Electronic band structure of graphene

Crystal lattice:



Electronic bands:



Linearity of bands around K points:


$$E(\mathbf{k}) \approx \pm \hbar v_F |\mathbf{k}| = \pm v_F |\mathbf{p}|$$

Electrons in graphene =
charged massless (relativistic) particles

$$E^2 = p^2 c^2 + m_0^2 c^4$$

Highlights of graphene physics

“Half-integer” and fractional QHE:





K. Novoselov et al., Nature 438, 197 (2005)
 Y. Zhang et al., Nature 438, 201 (2005)
 K. I. Bolotin et al., Nature 462, 196 (2009)

Room temperature QHE:

K. Novoselov et al., Science 315, 1379 (2007)

Universal dynamic conductivity:




R. R. Nair et al., Science 320, 1308 (2008)
 A. B. Kuzmenko et al., Phys. Rev. Lett. 100, 117401 (2008)



Other 2D Dirac-type materials

Silicene, germanene, phosphorene (?)

S. Cahangirov et al., Phys. Rev. Lett. 102, 236804 (2009)
 H. Liu et al., ACS Nano 8, 4033 (2014)





Artificial graphene, molecular graphene

C.-H. Park et al., PRL 101, 126804 (2008)
 M. Gibertini et al., Phys. Rev. B 79, 241406 (2009)
 L. Nádvorník et al., New J. Phys. 14, 053002 (2012)
 K. K. Gomes et al., Nature 483, 306 (2012)

Topological insulators (HgTe QWs, Bi_2Se_3 , Bi_2Te_3 , $\text{Bi}_x\text{Sb}_{1-x}$ etc.)

M. König et al., Science 318, 776 (2007)
 D. Hsieh et al., Nature 452, 970 (2008)
 H. Zhang et al., Nature Phys. 5, 438 (2009)



Semiconductors with a giant Rashba-type spin splitting (BiTeX , X = I, Cl or Br)

K. Ishizaka et al., Nature Materials 10, 521 (2011)
 S. Bordacs et al., Phys. Rev. Lett. 111, 166403 (2013)

Massless particles in 3D? Dirac and Weyl semimetals...

3D Dirac semimetals
(with spin-degenerate 3D conical bands)

Na₃Bi, Cd₃As₂

-
- Z. K Liu et al., Science 343, 864 (2014)
 - Z. K. Liu et al., Nature Mater. 13, 677 (2014)
 - S. Jeon et al., Nature Mater. 13, 851 (2014)
 - S. Borisenko et al., Phys. Rev. Lett. 113, 027603 (2014)
 - M. Neupane et al., Nature Comm. 5, 3786 (2014)
 - R. Y. Chen et al., Phys. Rev. Lett. 115, 176404 (2015)

3D Weyl semimetals
(3D conical bands with no spin degeneracy)

TaAs, NbAs

-
- B. Q. Lv et al., Nature Phys. 11, 724 (2015)
 - L. X. Yang et al., Nature Phys. 11, 728 (2015)
 - S.-Y. Xu et al., Nature Phys. 11, 748 (2015)



Dirac semimetal Cd_3As_2 = stable 3D analogue of graphene

nature
materials


LETTERS

PUBLISHED ONLINE: 25 MAY 2014 | DOI: 10.1038/NMAT3990


A stable three-dimensional topological Dirac semimetal Cd_3As_2

Z. K. Liu^{1†}, J. Jiang^{2,3†}, B. Zhou^{2,4†}, Z. J. Wang^{5†}, Y. Zhang¹⁴, H. M. Weng⁵, D. Prabhakaran², S-K. Mo⁴, H. Peng², P. Dudin⁶, T. Kim⁶, M. Hoesch⁶, Z. Fang⁵, X. Dai⁵, Z. X. Shen¹, D. L. Feng³, Z. Hussain⁴ and Y. L. Chen^{1,2,4,6*}

ARPES:



3D Dirac semimetal Cd_3As_2



-
- Z. K. Liu et al., Nature Mater. 13, 677 (2014)
 S. Jeon et al., Nature Mater. 13, 851 (2014)
 S. Borisenko et al., Phys. Rev. Lett. 113, 027603 (2014)
 M. Neupane et al., Nature Comm. 5, 3786 (2014)



Dirac semimetal Cd_3As_2 = stable 3D analogue of graphene

nature
materials


LETTERS

PUBLISHED ONLINE: 25 MAY 2014 | DOI: 10.1038/NMAT3990


A stable three-dimensional topological Dirac semimetal Cd_3As_2

Z. K. Liu^{1†}, J. Jiang^{2,3†}, B. Zhou^{2,4†}, Z. J. Wang^{5†}, Y. Zhang¹⁴, H. M. Weng⁵, D. Prabhakaran², S-K. Mo⁴, H. Peng², P. Dudin⁶, T. Kim⁶, M. Hoesch⁶, Z. Fang⁵, X. Dai⁵, Z. X. Shen¹, D. L. Feng³, Z. Hussain⁴ and Y. L. Chen^{1,2,4,6*}

ARPES:




3D Dirac semimetal Cd_3As_2



...but really not in line with other studies:

-
- I. Rosenman, J. Phys. Chem. Solids 30, 1385 (1969)
 - M. J. Aubin, et al., Phys. Rev. B 23, 3602 (1981)
 - H. Schleijpen et al., Int. J. Infrared Milli. 5, 171 (1984)
 - S. Jeon et al., Nature Mater. 13, 851 (2014)


Cd₃As₂ – Infrared reflectance at B=0



...in collaboration with C. C. Homes (Brookhaven)

Cd_3As_2 – Infrared reflectance at B=0

Two different crystallographic orientations of **tetragonal** Cd_3As_2



Cd₃As₂ – Infrared reflectance at B=0

Absorption of light in solids
(e.g., Fermi's golden rule):

$$\sigma_1(\omega) \propto \frac{\mathcal{D}(\omega)}{\omega}$$


For conical bands in 3D:

$$\mathcal{D}(\omega) \propto \omega^2$$

Absorption coefficient (optical conductivity) linear in photon frequency!

joint density
of states




Fairly high isotropy



see also T. Timusk et al., Phys. Rev. B 87, 235121 (2013)
M. Orlita et al., Nature Phys. 10, 233 (2014)



Absorption of light in graphene



"Flat" absorption of light (2.3%) defined only by the fine structure constant:




$$\alpha = \frac{e^2}{4\pi\epsilon_0\hbar c} \doteq \frac{1}{137}$$

R. R. Nair et al., Science 320, 1308 (2008)

A. B. Kuzmenko et al., Phys. Rev. Lett. 100, 117401 (2008)



Absorption of light in graphene



**Absorption of light in solids
(Fermi's golden rule):**

$$\lambda(\omega) \propto \frac{\mathcal{D}(\omega)}{\omega}$$

For conical bands in 2D:

$$\mathcal{D}(\omega) \propto \omega$$

Dispersionless (interband) absorption of light...

Cd₃As₂ – Optical response at B=0

Absorption of light in solids
(e.g., Fermi's golden rule):


$$\sigma_1(\omega) \propto \frac{\mathcal{D}(\omega)}{\omega}$$

joint density
of states

For conical bands in 3D:

$$\mathcal{D}(\omega) \propto \omega^2$$


Absorption coefficient linear in
photon frequency!





see also T. Timusk et al., Phys. Rev. B 87, 235121 (2013)
M. Orlita et al., Nature Phys. 10, 233 (2014)

Cd_3As_2 – Infrared reflectance at B=0

Two different crystallographic orientations of **tetragonal** Cd_3As_2




Cd₃As₂ – Infrared reflectance at B=0




Onsets of interband absorption
(Pauli blocking)


Cd₃As₂ – High-field magneto-reflectivity



Cd₃As₂ – High-field magneto-reflectivity




Cd_3As_2 – High-field magneto-reflectivity



Magneto-optical response linear in \sqrt{B} = typical signature of massless particles

Magneto-transmission of (multilayer epitaxial) graphene

Interband inter-Landau level transitions



Energy spectrum:

$$E_n = \pm v \sqrt{2e\hbar B |n|}$$


Velocity:

$$v = 1.02 \times 10^6 \text{ m/s}$$

Selection rules:


$$|n| \rightarrow |n| \pm 1$$

Cd_3As_2 – High-field magneto-reflectivity



Magneto-optical response linear in \sqrt{B} = typical signature of massless particles



Cd_3As_2 – High-field magneto-reflectivity



Magneto-optical response linear in \sqrt{B} = typical signature of massless particles

....massless yes, but not 3D Dirac


Dirac electrons – Landau level spectrum



$$E_n = \pm v \sqrt{2e\hbar B n + \hbar^2 k^2}$$

$$E_n = \pm v \sqrt{2e\hbar B n}$$

Cd_3As_2 – High-field magneto-reflectivity




Magneto-optical response linear in \sqrt{B} = typical signature of massless particles

....massless yes, but not 3D Dirac



Cd_3As_2 – Electronic band structure

3D Dirac semimetal Cd_3As_2



How the band structure looks like?

Where are the Dirac cones (if any)?

Energy scale of Dirac cones?

Some other conical bands?


Cd_3As_2 – Crystallographic structure

Tetragonal lattice with a large unit cell (~160 atoms)

G. Steigmann and J. Goodyear, Acta Crystallogr. B 24, 1062 (1968)
M. N. Ali et al. Inorganic Chemistry 53, 4062 (2014)

Ab initio calculations difficult for such a huge cell...

Z. Wang et al. Phys. Rev. B 88, 125427 (2013)



Cd_3As_2 – Crystallographic structure


Tetragonal lattice with a large unit cell (~ 160 atoms)

G. Steigmann and J. Goodyear, Acta Crystallogr. B 24, 1062 (1968)
M. N. Ali et al. Inorganic Chemistry 53, 4062 (2014)

Ab initio calculations difficult for such a huge cell...

Z. Wang et al. Phys. Rev. B 88, 125427 (2013)

Composed from distorted cubic sub-cells
= “almost” zinc-blende lattice





Bodnar model of electronic bands in Cd₃As₂

BAND STRUCTURE OF Cd₃As₂ FROM SHUBNIKOV-de HAAS
AND de HAAS - van ALPHEN EFFECTS


J. BODNAR

Department of Solid State Physics, Polish Academy of Sciences,
Zabrze, Poland

Experimental values of SdH and dHvA periods
and cyclotron effective masses found by
Rosenman and Doi et al. have been compared
with the theoretical predictions derived in
this work for a tetragonal narrow gap semi-
conductor. By the least square fit method
the values of band parameters were obtained.
It has been established that Cd₃As₂ has in-
verted band structure resembling HgTe under
tensile stress.

Bodnar model of electronic bands in Cd_3As_2

Standard Kane model for
zinc-blende semiconductors:




E. O. Kane, J. Phys. Chem. Solids 1, 249 (1957)

Valid for, e.g., GaAs, InAs, InSb, CdTe....



Bodnar model of electronic bands in Cd_3As_2

Kane model with a weak
tetragonal distortion:




J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw,(Elsevier, 1977) p. 311
see also H. Kildal, Phys. Rev. B 10, 5082 (1974).

...along tetragonal axis



Bodnar model of electronic bands in Cd_3As_2


Kane model with a weak
tetragonal distortion:



J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw, (Elsevier, 1977) p. 311

Bodnar model of electronic bands in Cd_3As_2

Kane model with a band **inversion** and
a weak **tetragonal distortion**:




J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw,(Elsevier, 1977) p. 311



Bodnar model of electronic bands in Cd₃As₂

Kane model with a band **inversion** and
a weak **tetragonal distortion**:





J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw,(Elsevier, 1977) p. 311



Bodnar model of electronic bands in Cd_3As_2

Standard Kane model
with a vanishing band gap:

Cube
(zinc-blende)




E. O. Kane, J. Phys. Chem. Solids 1, 249 (1957)

P. Kacman and W. Zawadzki, phys. stat. sol. (b) 47, 629 (1971)

M. Orlita et al., Nature Phys. 10, 233 (2014)

Massless Kane electrons in gapless HgCdTe



A single conical band in the center of Brillouin zone (due to accidental degeneracy of levels) hosting massless Kane electrons

In contrast to Dirac electrons, no protection against perturbations

E. O. Kane, J. Phys. Chem. Solids 1, 249 (1957)


P. Kacman and W. Zawadzki, phys. stat. sol. (b) 47, 629 (1971)

M. Orlita et al., Nature Phys. 10, 233 (2014)




Massless Kane electrons in gapless HgCdTe

Absorption coefficient linear
in photon energy:



Magneto-optical response
linear in \sqrt{B} :




Electronic bands in Cd_3As_2

Two kind of conical bands in Cd_3As_2 :

Two symmetry-protected **Dirac cones**
at the scale of crystal field splitting
($\delta < 50 \text{ meV}$)

A single cone of massless **Kane**
electrons, no symmetry protection
(energy scale $\sim 1 \text{ eV}$)

Γ point:



J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw, (Elsevier, 1977) p. 311

A. Akrap et al., submitted (2016)


Electronic bands in Cd_3As_2

Two kind of conical bands in Cd_3As_2 :

Two symmetry-protected **Dirac cones**
at the scale of crystal field splitting
($\delta < 50 \text{ meV}$)

A single cone of massless **Kane**
electrons, no symmetry protection
(energy scale $\sim 1 \text{ eV}$)

Γ point:





J. Bodnar, in Proc. III Conf. Narrow-Gap Semiconductors,
Warsaw, (Elsevier, 1977) p. 311


A. Akrap et al., submitted (2016)

3D massless Dirac and Kane electrons – Landau level spectrum


Landau levels - Dirac electrons



Landau levels - Kane electrons



Cd_3As_2 – High-field magneto-reflectivity



Magneto-optical signature of massless Kane electrons



Conclusions

Discovery/fabrication of graphene triggered vast search for other materials with analogous properties

Silicene, germanene, topological insulators, molecular graphene, artificial graphene, semiconductors with giant Rashba-type spin splitting...

Materials with 3D conical bands reported

Na_3Bi , Cd_3As_2 , zero-gap HgCdTe , ZrTe_5 ,
 TaAs , NbAs

-
- Z. K Liu et al., Science 343, 864 (2014)
M. Orlita et al., Nature Physics 10, 233 (2014)
Z. K. Liu et al., Nature Mater. 13, 677 (2014)
S. Jeon et al., Nature Mater. 13, 851 (2014)
R. Y. Chen et al., Phys. Rev. Lett. 115, 176404 (2015)
B. Q. Lv et al., Nature Phys. 11, 724 (2015)
L. X. Yang et al., Nature Phys. 11, 728 (2015)
S.-Y. Xu et al., Nature Phys. 11, 748 (2015)

Cd_3As_2 is most likely a symmetry-protected 3D Dirac semimetal

...but so far only massless Kane electrons have been seen experimentally

A. Akrap et al., submitted (2016)

Acknowledgements:

ERC-2012-AdG-320590 MOMB (M. Potemski)
Lia TeraMIR, TWINFUSYON